

1. For each pair of variables, write the equation (rearrange the equation if necessary) and determine the relationship. The relationships should be a description of how the first variable depends on the second.
 - a) area, A , and radius, r (for a circle)
 - b) volume, V , and radius, r (for a sphere)
 - c) displacement, d , and velocity, v (for an object in uniform motion)
 - d) displacement, d , and time, t (for an object accelerating uniformly from rest)
 - e) acceleration, a , and net force, F_{NET}
 - f) acceleration, a , and mass, m
 - g) force of friction, F_f , and normal force, F_N
 - h) displacement from equilibrium position, Δx , and spring constant, k
 - i) gravitational field strength, g , and mass, M
 - j) gravitational force, F_g , and separation distance, r
 - k) momentum, p , and velocity, v
 - l) gravitational potential energy, E_p , and height, h
 - m) kinetic energy, E_k , and mass, m
 - n) kinetic energy, E_k , and velocity, v
 - o) power, P , and time, t
 - p) current, I , and resistance, R

2. Determine the change in the following variables.
 - a) area, A , if radius, r , is doubled (for a circle)
 - b) volume, V , if radius, r , is halved (for a sphere)
 - c) displacement, d , if velocity, v , is increased by a factor of five (for an object in uniform motion)
 - d) displacement, d , if time, t , is decreased by a factor of four (for an object accelerating uniformly from rest)
 - e) acceleration, a , if net force, F_{NET} , is three times its original value
 - f) acceleration, a , if mass, m , is doubled
 - g) force of friction, F_f , if normal force, F_N , is one-third its original value
 - h) displacement from equilibrium position, Δx , if spring constant, k , is halved
 - i) gravitational field strength, g , if mass, M , is increased by a factor of four
 - j) gravitational force, F_g , if separation distance, r , is increased by a factor of six
 - k) momentum, p , if velocity, v is 2.5 times its original value
 - l) gravitational potential energy, E_p , if height, h , is decreased by a factor of ten
 - m) kinetic energy, E_k , if mass, m , is halved
 - n) kinetic energy, E_k , if velocity, v , is halved
 - o) power, P , if time, t , is twice its original value
 - p) current, I , if resistance, R , is $4/5$ its original value

3. Consider the equation for kinetic energy

$$E_k = \frac{1}{2}mv^2$$

where m represents the mass and v represents the velocity.

Determine the change in the kinetic energy for each of the following changes.

- The mass is doubled.
 - The velocity is doubled.
 - The mass is halved.
 - The velocity is halved.
 - The mass and velocity are both increased by a factor of three.
 - The mass and velocity are both decreased by a factor of three.
 - The mass is doubled and the velocity is halved.
 - The mass is decreased by a factor of four and the velocity is doubled.
4. Consider the equation for resistance

$$R = \frac{\rho \ell}{A}$$

where ρ represents the resistivity, ℓ represents the length of the wire, and A represents the cross-sectional area.

Determine the change in the resistance for each of the following changes.

- The length is increased by a factor of three.
 - The cross-sectional area is halved.
 - The length is decreased by a factor of four.
 - The cross-sectional area is increased by a factor of four.
 - The length and area are both doubled.
 - The length is tripled and the area is doubled.
 - The length is decreased by a factor of four and the area is tripled.
 - The length is decreased by a factor of six and the area is decreased by a factor of four.
5. Consider the equation for the gravitational force between two masses

$$F_g = G \frac{m_1 m_2}{r^2}$$

where G represents the gravitation constant, m_1 and m_2 represent the masses and r represents the separation distance.

Two objects are separated by a distance of 1000 km (from their centres). The gravitational force at this distance is 500 N. Determine the gravitational force between the masses for the following changes.

- One mass is doubled.
- Both masses are tripled.
- The distance separating the masses is increased to 5000 km.
- The distance separating the masses is decreased to 250 km.
- The distance separating the masses is increased to 1250 km.
- One mass is increased by a factor of five and the distance separating the masses is increased to 2000 km.
- Both masses are halved and the distance separating the masses is halved.
- One mass is tripled, the other is halved, and the distance separating them is decreased to 800 km.